



PolyStain TS Kit - for 1 Mouse and 2 Rabbit antibody on Rodent tissue

(DAB/Permanent Red/Ni-DAB)

NB-23-00137-1 (24mL)

NB-23-00137-2 (72mL)

NB-23-00137-3 (240mL)



**PolyStain TS Kit - for 1 Mouse and 2 Rabbit antibody on Rodent tissue
(DAB/Permanent Red/Ni-DAB)**

NB-23-00137-1; NB-23-00137-2; NB-23-00137-3

INTENDED USE:

Storage: 2-8°C

The PolyStain TS-MRR Kit is designed to use with user supplied one mouse and two rabbit primary antibodies to detect three distinct antigens on a single mouse/rat tissue or cell samples. This Kit has been tested on tissue specimens that are paraffin embedded. For frozen tissue a lower temperature of 65°C must be used for Antibody Blocker (Reagent 6) to prevent tissue from dissociating from slide. Please read through entire protocol as this protocol requires many step to be done in the defined order. Triple staining uses traditional methods in immunohistostaining to reveal three distinct antigens and their co-expression on a single tissue. PolyStain TS-MRR Kit from NeoBiotech Labs supplies polymer enzyme conjugates: polymer-HRP anti-mouse IgG, polymer HRP anti-rabbit IgG and polymer-AP anti-rabbit IgG with three substrates/chromogen; DAB (brown), Permanent Red (Red) and DAB-Ni (black). PolyStain TS-MRR Kit is a non-biotin system, avoiding non-specific binding caused by endogenous biotin. This kit has been optimized to have no cross detection when detecting more two primary antibodies from the same host species using unique blocking system. Simplified steps allow users to complete triple staining within 5 hours (without antigen retrieval) or 6-7 hours (with antigen retrieval). The well tested protocol provides user a method to permanently mount slides with coverslip.

KIT COMPONENTS:

Component No	Content	<u>24mL</u>	<u>72mL</u>	<u>240mL</u>
Reagent 1	Mouse Primer (RTU)	12mL	18mLx2	120mL
Reagent 2	Mouse HRP Polymer (RTU)	6mL	18mL	60mL
Reagent 3	Rabbit AP Polymer (RTU)	6mL	18mL	60mL
Reagent 4A	DAB Substrate (RTU)	15mL	18mLx2	120mL
Reagent 4B	DAB Chromogen (20x)	2mL	4mL	12mL
Reagent 5A	Permanent Red Substrate (RTU)	15mL	18mLx2	120mL
Reagent 5B	Permanent Red Activator (5x)	3mL	7.2mL	12mLx2
Reagent 5C	Permanent Red Chromogen (100x)	150µL	360µL	1.2mL
Reagent 6	Antibody Blocker (40x)	15mLx2	50mL	100mL
Reagent 7A	TS-MRR Blocker A (RTU)	12mL	18mLx2	120mL

Reagent 7B	TS-MRR Blocker B (RTU)	12mL	18mLx2	120mL
Reagent 8	Rabbit HRP Polymer (RTU)	12mL	18mLx2	120mL
Reagent 9A	DAB-Ni Substrate (20x)	1mL	2mL	6mL
Reagent 9B	Hydrogen Peroxide (20X)	1mL	2mL	6mL
Reagent 9C	Nickel Solution (7x)	3mL	6mL	18mL
Reagent 10	NeoMount Universal (RTU)	15mL	18mLx2	120mL

HRP = Horseradish Peroxidase AP = Alkaline Phosphatase Ms = Mouse Rb = Rabbit

PROTOCOL NOTES:

1. Proper Fixation: To ensure the quality of the staining and obtain reproducible performance, user needs to supply appropriately fixed tissue and well prepared slides.
2. Tissue needs to be adhered to the slide tightly to avoid falling off.
3. Paraffin embedded sections must be deparaffinize with xylene and rehydrated with a graded series of alcohols before staining.
4. Cell smear samples should be prepared as close to a monolayer as possible to obtain satisfactory results.
5. Control slides are recommended for interpretation of results: positive, reagent (slides treated with Isotype control reagent), and negative control.
6. **DO NOT** let specimen or tissue dry during protocol. This will generate false positive and/or false negative signal.
7. **Important:** Never combine two antibodies from the same host species in one incubation step. Incubate 1st primary mouse antibody with rabbit antibody.
8. The fixation, tissue section thickness, antigen retrieval and primary antibody dilution and incubation time effect results significantly. Investigator needs to consider all factors and determine optimal conditions when interpreting results.
9. We recommend TBS-T to be used as the wash buffer to get the highest sensitivity and clean background. Phosphate in the PBS-T may inhibit the activity of the alkaline phosphatase.

Note: 1X TBS-T =50mM Tris HCl, 150mM NaCl, 0.05% Tween-20 pH7.6.

NeoBiotech sells 10xTBS-T for your convenience (NB-23-00201)

Equipment or material needed but not provided:

1. Equipment and material for deparaffinization, such as fume absorbing hood, etc.
2. Heat source (microwave or hot plate) for HIER and antigen retrieval buffers
3. Thermometer
4. Timer, Beaker
5. Wash buffer: 0.01 M PBS with 0.5% Tween20, pH7.4
6. Peroxidase and alkaline phosphatase blocking buffer
7. 100% ethanol
8. 100% Xylene
9. Hematoxylin
10. Coverslip

Staining protocol selection and limitation of the kit:

- Most antigens will not be destroyed by heat. However, users need to check if there are proteins on the tissue that are heat sensitive before proceeding with the staining.
- NB-23-00137 Protocol-2 worksheet is suitable for one Mouse & one Rabbit primary Abs need pre-treatment, the other Rabbit primary Ab is sensitive to pre-treatment.
- NB-23-00137 Protocol-3 worksheet is suitable when one Mouse & one Rabbit primary antibody are sensitive to pre-treatment but the second Rabbit primary antibody needs pre-treatment.
- Please read the following table carefully before you start the experiment to ensure the result.
- This kit is not suitable for the following condition: 2 proteins are heat sensitive and detected by 2 rabbit antibodies and one mouse antibody requires HIER.

Staining protocol NB-23-00135 protocol-1:

Reagent	Staining Procedure	Incubation Time (Min.)
1. Peroxidase and Alkaline Phosphatase Blocking Reagent Not provided Fast, easy and it will block endogenous alkaline phosphatase	a. Incubate slides in PEROXIDASE BLOCKING REAGENT we recommend NeoPure Dual Enzyme Block NB-23-00193 b. Rinse the slide using distilled water.	10 min.

<p>2. Antigen retrieval (optional): Refer to primary antibody data sheet.</p>	<p>Note: Investigator needs to do antigen retrieval only one time during protocol see staining protocol.</p> <ol style="list-style-type: none"> Refer to primary antibody data sheet for antigen retrieval methods. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T (See note 9 above); 3 times for 2 minutes each. 	
<p>3. Primary Antibody Mix: Mix one Mouse and one Rabbit primary antibody Supplied by user.</p>	<p>Note: Investigator needs to optimize dilution prior to triple staining. DO NOT combine the same host species primary antibodies together at this step.</p> <ol style="list-style-type: none"> Apply 2 drops or enough volume of mouse and rabbit primary antibody mixture to cover the tissue completely. Incubate in moist chamber for 30- 60min. Recommend 30min to shorten total protocol time. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each. 	30 min
<p>4. Reagent 1 Mouse Primer (RTU)</p>	<ol style="list-style-type: none"> Apply 1 to 2 drops (50-100µL) of Reagent 1 (Mouse Primer) to cover the tissue completely. Incubate slides in moist chamber for 15 min. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each. 	10 min
<p>5. Mix Reagent 2: Mouse AP Polymer (RTU) with Reagent 3: Rabbit HRP Polymer (RTU)</p>	<p>Note: Make sufficient polymer mixture by adding Reagent 2 (Mouse AP Polymer) and Reagent 3 (Rabbit HRP Polymer) at 1:1 ratio, mix well. Do not mix more than you need for the experiment because the polymer mixture may not be as stable as non-mixed polymer.</p> <ol style="list-style-type: none"> Apply 1 to 2 drops (50-100µL) of the mixture to cover the tissue completely. Incubate in moist chamber for 30 min. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each 	30 min
<p>6. Reagent 4A&4B 4A: DAB Substrate(RTU) 4B: DAB Chromogen (20x)</p>	<p>Note: Make enough DAB mix by adding 1 drop of Reagent 4B (DAB Chromogen) in 1mL of Reagent 4A (DAB Substrate). Mix well. Use within 7 hours store at 4°C.</p> <ol style="list-style-type: none"> Apply 1 to 2 drops (50-100µL) of your DAB mixture to cover the tissue completely. Incubate for 5min. Rinse thoroughly with distilled water. Wash with 1xTBS-T only; 3 times for 2 minutes each. 	5 min
<p>7. Reagent 5A, 5B, 5C Reagent 5A: Permanent Red Substrate (RTU)</p>	<p>Note: Shake Permanent Red Activator before adding into Permanent Red Substrate.</p> <ol style="list-style-type: none"> Add 200µL of Reagent 5B (Activator) into 1mL of Reagent 5A (Substrate) and mix well. Add 10µL of Reagent 5C (Chromogen) into the mixture and mix well. [Note: For fewer slides, Add 100µL of Reagent 5B (Activator) into 500µL of Reagent 5A 	10 min

<p>Reagent 5B: Permanent Red Activator (5x)</p> <p>Reagent 5C: Permanent Red Chromogen (100x)</p> <p>To get maximum sensitivity of AP polymer, Please repeat chromogen step</p>	<p>(Substrate) and mix well. Add 5µL of Reagent 5C (Chromogen) into the mixture and mix well.]</p> <p>b. Apply 2 drops (100µL) or enough volume of Permanent Red working solution to completely cover the tissue. Incubate for 10 min, observe appropriate color development. To increase AP signal aspirate or tap off chromogen and apply 2-3 drops (100µL) again of the Permanent Red working solution to completely cover the tissue for additional 5 to 10min</p> <p>c. Rinse well with distilled water</p>	
<p>8. Reagent 6</p> <p>Antibody Blocker (40x)</p>	<p>Note: This step will block antibodies of previous step so no cross reaction will occur in this protocol. HIER can be done immediately after Antibody Blocker step if the primary antibodies requires antigen retrieval. For frozen tissues, a lower temperature of 65°C must be used during the Antibody Blocker step to prevent dissociation of the tissue from the slide.</p> <p>a. Use hot plate or water bath to heat diluted Reagent 6 (Antibody Blocker) to 1x solution (1 part of Antibody Blocker in 39 parts of distilled water) to 80°C. Make enough volume to cover the tissue in beaker.</p> <p>b. Put slides in heated Antibody Blocker for 10 minutes at 80°C.</p> <p>c. Remove slides from the Antibody blocker; cool slides 5 seconds.</p> <p>d. Rinse slides in multiple changes of distilled water. If antigen retrieval step is required go directly to step 9 if not complete step 8e and move on to step 10.</p> <p>e. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.</p>	<p>10 min</p>
<p>9. Antigen retrieval: Refer to primary antibody data sheet.</p>	<p>a. Refer to primary antibody data sheet for antigen retrieval methods.</p> <p>b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.</p>	<p>Up to 1 hour</p>
<p>10. Reagent 7A TS-MRR Blocker A (RTU)</p>	<p>a. Apply 2 drops or enough volume of Reagent 7A (DS-MRR Blocker A) to cover the tissue completely. Mix well on the slide and incubate in moist chamber for 30 min.</p> <p>b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each</p>	<p>30 min</p>
<p>11. Reagent 7B TS-MRR Blocker B (RTU)</p>	<p>a. Apply 2 drops or enough volume of Reagent 7B (DS-MRR Blocker B) to cover the tissue completely. Mix well on the slide and Incubate in moist chamber for 5 min.</p> <p>b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.</p>	<p>5 min</p>

<p>12. 2nd Mouse primary antibody</p> <p>Supplied by user.</p>	<p>Note: Investigator needs to optimize dilution prior to triple staining.</p> <p>a. Apply 2 drops or enough volume of the 2nd mouse primary antibody to cover the tissue completely. Incubate in moist chamber for 30-60 min. Recommend 30 minutes to shorten total protocol time.</p> <p>b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.</p>	<p>30 min</p>
<p>13. Reagent 8 Rabbit HRP Polymer (RTU)</p>	<p>a. Apply 1 to 2 drops (50-100µL) of Reagent 8 (Rabbit HRP Polymer) to cover the tissue completely. Incubate slides in moist chamber for 15 min.</p> <p>b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.</p>	<p>15 min</p>
<p>14. Reagent 9A, 4B, 9C&9C</p> <p>9A: DAB-Ni Substrate (20x)</p> <p>4B: DAB Chromogen (20x)</p> <p>9B: Hydrogen Peroxide (20x)</p> <p>9C: Nickel Solution (7x)</p>	<p>a. Prepare 1mL of distilled water. Add 1 drop of Reagent 9A (DAB-Ni Substrate) into 1mL of distilled water. Mix well.</p> <p>b. Add 1 drop of Reagent 4B (DAB Chromogen) and 1 drop of concentrated Reagent 9B (Hydrogen Peroxide) to the diluted Reagent. Mix well.</p> <p>c. Add 3 drops of Reagent 9C (Nickel Solution) to the mixture. Mix well.</p> <p>d. Add about 100µL (2 drops) of DAB-Ni working solution to each slide and incubate in an enclosed chamber at room temperature for about 5 minutes. When appropriate color is developed, rinse under tap water gently for about 1- 2 minutes.</p> <p>Note: Use DAB-Ni working solution within 7 hours and store at 4°C keeping away from light during operation.</p>	<p>5 min</p>
<p>15. HEMATOXYLIN Not provided</p>	<p>a. Counterstain with 2 drops (100µL) or enough volume of hematoxylin to completely cover tissue. Incubate for 10-15 seconds.</p> <p>b. Rinse thoroughly with tap water for 2-3min.</p> <p>c. Put slides in PBS until show blue color (about ½ - 1min.)</p> <p>d. Rinse well in distilled water</p>	<p>10 -15 sec</p>
<p>16. Reagent 10: NeoMount Universal (RTU)</p>	<p>a. Apply 2 drops (100µL) or enough volume of Reagent 10 (NeoMount Universal) to cover tissue when tissue is wet. Rotate the slides to allow NeoMount Universal spread evenly.</p> <p>b. Place slides horizontally in an oven at 40-50°C for at least 30 minutes or leave it at room temperature until slides are thoroughly dried.</p>	

Trouble shoot:

Problem	Tips
Uneven stain on 3 primary antibodies	1. Need to adjust the titer of each antibody. 2. The amount of each protein expressed on tissue may be different
No stain on 1 or 2 antibodies	Missing steps or step reversed.

PROTOCOL NOTES:

1. The fixation, tissue slide thickness, antigen retrieval and primary antibody dilution and incubation time affect results significantly. Investigator needs to consider all factors and determine optimal conditions when interpreting the result.
2. Permanent Red is insoluble in organic solvent and can be coverslipped as well. However the dehydration steps must be shorter for optimal tissue structure and chromogen signal maintenance.

Note: Please wipe off extra water and air dry slides before dehydration and clear.

- a. 1x 80% Ethanol 20 seconds;
- b. 1x 95% Ethanol 20 seconds;
- c. 3x 100% Ethanol 20 seconds each;
- d. 1x 100% Xylene 20 seconds;
- e. Add 1 drop of xylene based mountant (Cat. No. NeoMount Perm, NB-23-00156) and coverslip. Press to push the air bubble out.

CAUTION: DO NOT dehydrate in xylene longer than 20 seconds! It will erase Permanent Red stain!

PRECAUTIONS:

Please wear gloves, eye protection and take other necessary precautions. If any of the reagent come in contact with skin wash area completely with plenty of water and soap. If irritation develops seek medical attention.

For research use only.

Work Sheet for NB-23-00137 Kit

We designed this work sheet to help you track of each step. We recommend you use this sheet to record the actual time of each step conducted as it will be helpful for questions with our technical support. To insure that all steps are done properly, we recommend that the user fill in the actual time of their experimental step and any variation. Results will vary if time recommendations are not followed. RTU translates to ready to use.

- Used for tester to check “√” each step during the experiment
- Steps follow after de-paraffinization
- Refer to insert for details of each step

NB-23-00137 Protocol-1 is suitable when all primary antibodies need pre-treatment or all primary antibodies do not need pre-treatment

	Main Protocol Step	NB-23-00137 Protocol-1	Experiment 1 Date:	Experiment 2 Date:	Experiment 3 Date:	Experiment 4 Date:
1	Step 1	Peroxidase & Alkaline Phosphatase Block NB-23-00193 is recommended. User supplied				
2	Step 2	HIER(Optional)				
3	Step 3	Mouse 1°Ab & Rabbit 1°Ab mix User supplied (30-60min)				
4	Step 4	Reagent 1 Mouse primer RTU 15min				
5	Step 5	Reagent 2&Reagent 3 Mouse AP Polymer & Rabbit HRP Polymer require mixing (30min)				
6	Step 6	Reagent 4A& Reagent 4B DAB requires mixing. (5min) Wash with 1xTBS-T after rinse well with distilled water				
7	Step 7	Reagent 5A, Reagent 5B Reagent 5C Permanent Red requires mixing. (10min)				

8	Step 8	Reagent 6 Antibody Blocker requires mixing. (10min)				
9	Step 10	Reagent 7A DS-MRR Blocker A RTU (30min)				
10	Step 11	Reagent 7B DS-MRR Blocker B RTU (5min)				
11	Step 12	Mouse 1°Ab User supplied (30-60 min)				
12	Step 13	Reagent 8 Rabbit HRP Polymer RTU (15 min)				
13	Step 14	Reagent 9A,9B,9C&4B DAB-Ni requires mixing (5min)				
14	Step 15	Counter stain User supplied				
15	Step 16	Reagent 10 NeoMount Universal RTU				
16	Result	Stain pattern on controls are correct: Fill in Yes or NO				

Note: 1.Normal wash steps = Wash with PBS-T containing 0.05% Tween-20 or **1X TBS-T**; 3 times for 2 minutes each.

Testing result:

NB-23-00137 Protocol-2 is suitable when one Mouse & one Rabbit primary antibodies need pre-treatment, but the second Mouse primary antibodies is sensitive to pre-treatment.

	Main Protocol Step	NB-23-00137 Protocol-2	Experiment 1 Date:	Experiment 2 Date:	Experiment 3 Date:	Experiment 4 Date:
1	Step 1	Peroxidase or Alkaline Phosphatase Block NB-23-00193 is recommended. User supplied				
2	Step 12	Mouse 1°Ab (sensitive to HIER) User supplied (30-60min)				
3	Step 13	Reagent 8 (RTU) Rabbit HRP Polymer RTU (15min)				
4	Step 6	Reagent 4A&4B DAB requires mixing (5 min)				
5	Step 8	Reagent 6 Antibody Blocker requires mixing (10min)				
6	Step 9	HIER (DAB will not be removed)				
7	Step 10	Reagent 7A (RTU) DS-MRR Blocker A RTU (30min)				
8	Step 11	Reagent 7B (RTU) DS-MMR Blocker B RTU (5min)				
9	Step 3	Mouse 1°Ab & Rabbit 1°Ab mix (Abs requires HIER) User supplied (30-60 min)				

10	Step 4	Reagent 1 Mouse primer RTU 15min				
11	Step 5	Reagent 2&Reagent 3 Mouse AP Polymer & Rabbit HRP Polymer require mixing (30min) Wash with 1x TBS-T				
12	Step 7	Reagent 5A, Reagent 5B & Reagent 5C Permanent Red requires mixing. (10min)				
13	Step 14	Reagent 9A,9B,9C&4B DAB-Ni requires mixing (5min)				
14	Step 15	Counter stain User supplied				
15	Step 16	Reagent 10 NeoMount Universal RTU				
16	Result	Stain pattern on controls are correct: Fill in Yes or NO				

Note1: Normal wash steps = Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.

Testing result:

NB-23-00137 Protocol-3 is suitable when one Mouse & one Rabbit primary antibodies are sensitive to pre-treatment but the second Mouse primary antibody needs pre-treatment.

	Main Protocol Step	NB-23-00137 Protocol-3	Experiment 1 Date:	Experiment 2 Date:	Experiment 3 Date:	Experiment 4 Date:
1	Step 1	Peroxidase or Alkaline Phosphatase Block NB-23-00193 is recommended. User supplied				
2	Step 3	Mouse 1°Ab & Rabbit 1°Ab mix User supplied (30-60min.)				
3	Step 4	Reagent 1 Mouse primer RTU 15min				
4	Step 5	Reagent 2&Reagent 3 Mouse HRP Polymer & Rabbit AP Polymer require mixing. (30min)				
5	Step 6	Reagent 3A&Reagent 3B DAB require mixing. (5min) Wash with 1xTBS-T after rinse well with distilled water				
6	Step 7	Reagent 5A, Reagent 5B &Reagent 5C Permanent Red requires mixing. (10min)				
7	Step 8	Reagent 6 Antibody Blocker required mixing. (10min)				
8	Step 9	HIER Refer to antibody datasheet.				
9	Step 10	Reagent 7A DS-MRR Blocker A RTU (30min)				

10	Step 11	Reagent 7B DS-MRR Blocker B RTU (5min)				
11	Step 12	Rabbit 1°Ab (Not sensitive to HIER) User supplied (30-60min.)				
12	Step 13	Reagent 8 Rabbit HRP Polymer (RTU) (15min.)				
13	Step 14	Reagent 9A,9B,9C&4B DAB-Ni requires mixing (5min)				
14	Step 15	Counter stain User supplied				
15	Step 16	Reagent 10 NeoMount Universal RTU				
16	Result	Stain pattern on controls are correct: Fill in Yes or NO				

Note1: Normal wash steps = Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T; 3 times for 2 minutes each.

Testing result:

